

REIMBURSEMENT, BILLING, AND CODING GUIDE FOR TRODELVY® (sacituzumab govitecan-hziy)

The tables below include examples of codes that may be appropriate for use when billing and seeking reimbursement for treatment with TRODELVY.

Coding requirements may vary by insurer or plan. Gilead Sciences, Inc. has provided these codes only as a reference. When submitting a claim for TRODELVY, always verify coding requirements with the relevant payer. Healthcare professionals are solely responsible for selecting codes that appropriately reflect the patient's diagnosis, the services rendered, and the applicable payer's guidelines. The use of this information does not guarantee payment or that any payment received will cover costs.

INDICATIONS

TRODELVY® (sacituzumab govitecan-hziy) is a Trop-2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with:

- Unresectable locally advanced or metastatic triple-negative breast cancer (mTNBC) who have received two or more prior systemic therapies, at least one of them for metastatic disease.
- Locally advanced or metastatic urothelial cancer (mUC) who have previously received a platinum-containing chemotherapy and either programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor. This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

Please see Important Safety Information, including BOXED WARNINGS on Neutropenia and Diarrhea below.

HEALTHCARE COMMON PROCEDURE CODING SYSTEM (HCPCS)

HCPCS coding requirements will vary by payer, setting of care, and date of service. Please verify patient-specific insurance benefits to confirm specific coding and billing guidelines for TRODELVY.

HCPCS Level II Code	Description	Note
J9317 (effective January 1, 2021)	Injection, sacituzumab govitecan-hziy, 2.5 mg	For all providers and settings of care for which HCPCS codes are reported

As of January 1, 2017, Medicare claims require the use of the JW modifier (drug amount discarded/not administered to any patient) when applicable. Other payers may have similar requirements.

NATIONAL DRUG CODE (NDC)

Payer requirements regarding the use of a 10-digit or 11-digit NDC may vary. Both formats are listed here for your reference. Please consult with the payer to understand specific billing requirements.

NDC	Code	Description
10-digit code	55135-132-01	TRODELVY for injection is supplied as 180 mg of sacituzumab govitecan-hziy as lyophilized powder in a single-use vial
11-digit code (with leading 0)	55135-0132-01	

CURRENT PROCEDURAL TERMINOLOGY (CPT®) CODE FOR DRUG ADMINISTRATION SERVICE

The recommended dose of TRODELVY is administered as an intravenous infusion once weekly on Days 1 and 8 of continuous 21-day treatment cycles. Please refer to the Full Prescribing Information for complete dosing and administration guidelines.

CPT Code	Description
96413	Chemotherapy administration, intravenous infusion technique; up to 1 hour, single or initial substance/drug
96415	Chemotherapy administration, intravenous infusion technique; each additional hour (list separately in addition to code for primary procedure)

CPT Copyright 2019 American Medical Association. All rights reserved. CPT® is a registered trademark of the American Medical Association.

IMPORTANT SAFETY INFORMATION

BOXED WARNING: NEUTROPENIA AND DIARRHEA

- **Severe or life-threatening neutropenia may occur. Withhold TRODELVY for absolute neutrophil count below 1500/mm³ or neutropenic fever. Monitor blood cell counts periodically during treatment. Consider granulocyte colony-stimulating factor (G-CSF) for secondary prophylaxis. Initiate anti-infective treatment in patients with febrile neutropenia without delay.**
- **Severe diarrhea may occur. Monitor patients with diarrhea and give fluid and electrolytes as needed. Administer atropine, if not contraindicated, for early diarrhea of any severity. At the onset of late diarrhea, evaluate for infectious causes and, if negative, promptly initiate loperamide. If severe diarrhea occurs, withhold TRODELVY until resolved to ≤Grade 1 and reduce subsequent doses.**

Please see accompanying full [Prescribing Information](#), including **BOXED WARNING**, and continued **Important Safety Information** on the next page.



REVENUE CODES (FOR HOSPITAL CLAIMS ONLY)

All hospital claim forms must include a revenue code for each charge line item. The following revenue codes are most relevant for physician-administered drugs.

Revenue Code	Description
0250	Pharmacy
0636	Pharmacy—drugs requiring detailed coding

ICD-10-CM DIAGNOSIS CODES

ICD-10 diagnosis codes represent medical terminology for diseases, disorders, or other medical conditions affecting the patient. Proper diagnosis coding involves using the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) volumes to select the appropriate codes based on documentation in the patient's medical record and assigning those codes correctly on claims.

ICD-10-CM Diagnosis Code	Description
C50.212* (example)	Malignant neoplasm of upper-inner quadrant of left female breast
C50.411* (example)	Malignant neoplasm of upper-outer quadrant of right female breast
C65.* (example)	Malignant neoplasm of renal pelvis
C66.* (example)	Malignant neoplasm of ureter
C67.* (example)	Malignant neoplasm of bladder
C68.* (example)	Malignant neoplasm of other and unspecified urinary organs

*These are only 6 examples of applicable ICD-10-CM codes. To see all ICD-10-CM codes for breast cancer and bladder cancer see https://www.cms.gov/icd10m/version37-fullcode-cms/fullcode_cms/P0230.html and https://www.cms.gov/icd10m/version37-fullcode-cms/fullcode_cms/P0245.html.

Gilead Sciences, Inc. is not responsible for the accuracy of this site.

IMPORTANT SAFETY INFORMATION (CONTINUED)

CONTRAINDICATIONS

- Severe hypersensitivity to TRODELVY.

WARNINGS AND PRECAUTIONS

Neutropenia: Severe, life-threatening, or fatal neutropenia can occur and may require dose modification. Neutropenia occurred in 61% of patients treated with TRODELVY. Grade 3-4 neutropenia occurred in 47% of patients. Febrile neutropenia occurred in 7%. Withhold TRODELVY for absolute neutrophil count below 1500/mm³ on Day 1 of any cycle or neutrophil count below 1000/mm³ on Day 8 of any cycle. Withhold TRODELVY for neutropenic fever.

Diarrhea: Diarrhea occurred in 65% of all patients treated with TRODELVY. Grade 3-4 diarrhea occurred in 12% of patients. One patient had intestinal perforation following diarrhea. Neutropenic colitis occurred in 0.5% of patients. Withhold TRODELVY for Grade 3-4 diarrhea and resume when resolved to ≤Grade 1. At onset, evaluate for infectious causes and if negative, promptly initiate loperamide, 4 mg initially followed by 2 mg with every episode of diarrhea for a maximum of 16 mg daily. Discontinue loperamide 12 hours after diarrhea resolves. Additional supportive measures (e.g., fluid and electrolyte substitution) may also be employed as clinically indicated. Patients who exhibit an excessive cholinergic response to treatment can receive appropriate premedication (e.g., atropine) for subsequent treatments.

Please see accompanying full [Prescribing Information](#), including **BOXED WARNING**, and continued Important Safety Information on the next page.

 **TRODELVY**[™]
sacituzumab govitecan-hziy
180 mg for injection

IMPORTANT SAFETY INFORMATION (CONTINUED)

WARNINGS AND PRECAUTIONS (cont'd)

Hypersensitivity and Infusion-Related Reactions:

Serious hypersensitivity reactions including life-threatening anaphylactic reactions have occurred with TRODELVY. Severe signs and symptoms included cardiac arrest, hypotension, wheezing, angioedema, swelling, pneumonitis, and skin reactions. Hypersensitivity reactions within 24 hours of dosing occurred in 37% of patients. Grade 3-4 hypersensitivity occurred in 2% of patients. The incidence of hypersensitivity reactions leading to permanent discontinuation of TRODELVY was 0.3%. The incidence of anaphylactic reactions was 0.3%. Pre-infusion medication is recommended. Observe patients closely for hypersensitivity and infusion-related reactions during each infusion and for at least 30 minutes after completion of each infusion. Medication to treat such reactions, as well as emergency equipment, should be available for immediate use. Permanently discontinue TRODELVY for Grade 4 infusion-related reactions.

Nausea and Vomiting: Nausea occurred in 66% of all patients treated with TRODELVY and Grade 3 nausea occurred in 4% of these patients. Vomiting occurred in 39% of patients and Grade 3-4 vomiting occurred in 3% of these patients. Premedicate with a two or three drug combination regimen (e.g., dexamethasone with either a 5-HT₃ receptor antagonist or an NK₁ receptor antagonist as well as other drugs as indicated) for prevention of chemotherapy-induced nausea and vomiting (CINV). Withhold TRODELVY doses for Grade 3 nausea or Grade 3-4 vomiting and resume with additional supportive measures when resolved to Grade ≤ 1 . Additional antiemetics and other supportive measures may also be employed as clinically indicated. All patients should be given take-home medications with clear instructions for prevention and treatment of nausea and vomiting.

Increased Risk of Adverse Reactions in Patients

with Reduced UGT1A1 Activity: Patients homozygous for the uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1)*28 allele are at increased risk for neutropenia, febrile neutropenia, and anemia and may be at increased risk for other adverse reactions with TRODELVY. The incidence of Grade 3-4 neutropenia was 67% in patients homozygous for the UGT1A1*28, 46% in patients heterozygous for the UGT1A1*28 allele and 46% in patients homozygous for the wild-type allele. The incidence of Grade 3-4 anemia was 25% in patients homozygous for the UGT1A1*28 allele, 10% in patients heterozygous for the UGT1A1*28 allele, and 11% in patients homozygous for the wild-type allele. Closely monitor patients with known reduced UGT1A1 activity for adverse reactions. Withhold or permanently discontinue TRODELVY based on clinical

assessment of the onset, duration and severity of the observed adverse reactions in patients with evidence of acute early-onset or unusually severe adverse reactions, which may indicate reduced UGT1A1 function.

Embryo-Fetal Toxicity: Based on its mechanism of action, TRODELVY can cause teratogenicity and/or embryo-fetal lethality when administered to a pregnant woman. TRODELVY contains a genotoxic component, SN-38, and targets rapidly dividing cells. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TRODELVY and for 6 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with TRODELVY and for 3 months after the last dose.

ADVERSE REACTIONS

In the ASCENT study (IMMU-132-05), the most common adverse reactions (incidence $\geq 25\%$) were fatigue, neutropenia, diarrhea, nausea, alopecia, anemia, constipation, vomiting, abdominal pain, and decreased appetite. The most frequent serious adverse reactions (SAR) ($>1\%$) were neutropenia (7%), diarrhea (4%), and pneumonia (3%). SAR were reported in 27% of patients, and 5% discontinued therapy due to adverse reactions. The most common Grade 3-4 lab abnormalities (incidence $\geq 25\%$) in the ASCENT study were reduced neutrophils, leukocytes, and lymphocytes.

In the TROPHY study (IMMU-132-06), the most common adverse reactions (incidence $\geq 25\%$) were diarrhea, fatigue, neutropenia, nausea, any infection, alopecia, anemia, decreased appetite, constipation, vomiting, abdominal pain, and rash. The most frequent serious adverse reactions (SAR) ($\geq 5\%$) were infection (18%), neutropenia (12%, including febrile neutropenia in 10%), acute kidney injury (6%), urinary tract infection (6%), and sepsis or bacteremia (5%). SAR were reported in 44% of patients, and 10% discontinued due to adverse reactions. The most common Grade 3-4 lab abnormalities (incidence $\geq 25\%$) in the TROPHY study were reduced neutrophils, leukocytes, and lymphocytes.

Please see accompanying full [Prescribing Information](#), including **BOXED WARNING**, and continued Important Safety Information on the next page.

 TRODELVY™
sacituzumab govitecan-hziy
180 mg for injection

IMPORTANT SAFETY INFORMATION (CONTINUED)

DRUG INTERACTIONS

UGT1A1 Inhibitors: Concomitant administration of TRODELVY with inhibitors of UGT1A1 may increase the incidence of adverse reactions due to potential increase in systemic exposure to SN-38. Avoid administering UGT1A1 inhibitors with TRODELVY.

UGT1A1 Inducers: Exposure to SN-38 may be substantially reduced in patients concomitantly receiving UGT1A1 enzyme inducers. Avoid administering UGT1A1 inducers with TRODELVY.

Please see accompanying full [Prescribing Information](#), including **BOXED WARNING**, also at TRODELVY.com/prescribing-information.

For more information about reimbursement, billing, and coding for TRODELVY, please contact:

TRODELVY ACCESS SUPPORT

Phone: 1-844-TRODELVY (876-3358) Monday – Friday, 9 AM - 7 PM ET

www.TRODELVY.com



GILEAD

TRODELVY, the TRODELVY Logo, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies.

©2021 Gilead Sciences, Inc. All rights reserved.

2021-US-TROT-00107 03/21



TRODELVY™

sacituzumab govitecan-hziy

180 mg for injection